

## Selected References

- Ada L, Dean CM, Hall JM, et al. A treadmill and overground walking program improves walking in persons residing in the community after stroke: a placebo-controlled randomized trial. *Arch Phys Med Rehabil.* 2003;84:1486-1491.
- Alexander NB, Schultz AB, Warwick DN. Rising from a chair: effects of age and functional ability on performance biomechanics. *J Gerontol Med Sci.* 1991;46:M91-98.
- American College of Sports Medicine, ed. *ACSM's Guidelines for Exercise Testing and Prescription.* 8<sup>th</sup> ed. Baltimore MD: American College of Sports Medicine; 2010.
- American College of Sports Medicine, Chodzko-Zajko WJ, Proctor DN, et al. American college of sports medicine position stand. Exercise and physical activity for older adults. *Med Sci Sport Exerc.* 2009;41:1510-1530.
- APTA White Paper – The Human Movement System. August, 2015.
- Avers D, Brown M. White paper: strength training for the older adult. *J Geri Phys Ther.* 2009;32(4):148-152.
- Barbieri G, Gissot A-S, Fouque F et al. Does proprioception contribute to the sense of verticality? *Exp Brain Res.* 2008;185:545-552.
- Barbieri G, Gissot A-S, Perennou D. Ageing of the postural vertical. *Age.* 2010;32:51-60.
- Bastian AJ, Martin TA, Keating JG, Thach WT. Cerebellar ataxia: abnormal control of interaction torques across multiple joints. *J Neurophysiology.* 1996;76:492-509.
- Bastian AJ. Mechanisms of ataxia. *Phys Ther.* 1997;77:672-675.
- Bastian AJ, Zackowski KM, Thach WT. Cerebellar ataxia: torque deficiency or torque mismatch between joints? *J Neurophysiology.* 2000;83:3019-3030.
- Bastian AJ. Cerebellar limb ataxia: abnormal control of self-generated and external forces. *Ann NY Acad Sci.* 2002;978:16-27.
- Bastian AJ. Learning to predict the future: the cerebellum adapts feedforward movement control. *Curr Opin Neurobiol.* 2006;16:645-649.
- Bastian AJ. Understanding sensorimotor adaptation and learning for rehabilitation. *Current Opinion Neurol.* 2008;21:628-633.
- Bogey R and Hornby TG. Gait training strategies utilized in poststroke rehabilitation: are we really making a difference? *Top Stroke Rehabil.* 2007;14:1-8.
- Behrman AL, Ardolina E, Vanhiel LR, et al. Assessment of functional improvement without compensation reduces variability of outcome measures after human spinal cord injury. *Arch Phys Med Rehabil.* 2012;93(9):1518-1529.
- Behrman AL, Teitelbaum P, Cauraugh JH. Verbal instructional sets to normalize the temporal and spatial gait variables in Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 1998;65:580-582.
- Behrman AL, Cauraugh JH, Light KE. Practice as an intervention to improve speeded motor performance and motor learning in Parkinson's disease. *J Neurolog Sci.* 2000;174:127-136.
- Birkenmeier RL, Prager EM, Lang CE. Translating animal doses of task-specific training to people with chronic stroke in 1-hour therapy sessions: a proof-of-concept study. *Neurorehabil Neural Repair.* 2010;24(7):620-630.

Bland MD, Whitson M, Harris H, et al. Descriptive data analysis examining how standardized assessments are used to guide post-acute discharge recommendations for rehabilitation services after stroke. *Phys Ther*. 2015 May;95(5):710-9.

Bloem BR, de Vries NM, Ebersbach G. Nonpharmacological treatments for patients with Parkinson's disease. *Movement Disorders*. 2015. 30(11):1504-1520.

Burleigh-Jacobs A, Horak FB, Nutt JG, et al. Step initiation in Parkinson's disease: influence of Levodopa and external sensory triggers. *Movement Disorders*. 1997;12:206-215.

Carr J and Shepherd R. *Stroke Rehabilitation: guidelines for exercise and training to optimize motor skill*. London: Butterworth-Heinemann; 2003:230-231, 242-243.

Coffin-Zadai CA. Disabling our diagnostic dilemmas. *Phys Ther*. 2007;87:641-653.

Cunningham R, Iansek R, and Bradshaw JL. Movement-related potentials in Parkinson's disease: external cues and attentional strategies. *Movement Disorders*. 1999;14:63-68.

da Cunha Filho IT, Lim PAC, Qureshy H, Henson H, Monga T, Protas EJ. A comparison of regular rehabilitation and regular rehabilitation with supported treadmill ambulation training for acute stroke patients. *Journal of Rehabilitation Research and Development*. 2001;38.

Danielsson A, Sunnerhagen KS. Oxygen consumption during treadmill walking with and without body weight support in patients with hemiparesis after stroke and in healthy subjects. *Arch Phys Med Rehabil*. 2000;81:953-957.

Dam M, Tonin P, Casson S, et al. Effects of conventional and sensory-enhanced physiotherapy on disability of Parkinson's disease patients. *Adv Neurol*. 1996;69:551-555.

Dean CM, Richards CL, Malouin F. Task-related circuit training improves performance of locomotor tasks in chronic stroke: a randomized controlled pilot trial. *Arch Phys Med Rehabil*. 2000;81:409-417.

Dean CM and Shepherd RB. Task-related training improves performance of seated reaching tasks after stroke: a randomized controlled trial. *Stroke*. 1997;28:722-728.

DeJong G, Hsieh CH, Putman K, et al. Physical therapy activities in stroke, knee arthroplasty, and traumatic brain injury rehabilitation: their variation, similarities, and association with functional outcomes. *Phys Ther*. 2011;91(12):1826-37.

DeJong S., Schaefer SY, Lang CE. Need for speed: better movement quality during faster task performance after stroke. *Neurorehabil Neural Repair*. 2012;26(4):362-73.

Deschenes MR and Kraemer WJ. Performance and physiologic adaptations to resistance training. *Am J Phys Med Rehabil*. 2002;81(11 Suppl):S3-S16.

De Vreede PL, van Meeteren NL, Samson MM, et al. Functional tasks exercise versus resistance to improve daily function in older women: a feasibility study. *Arch Phys Med Rehab*. 2004;85:1952-1961.

Dibble LE, Nicholson DE, Shultz B, et al. Sensory cueing effects on maximal speed gait initiation in persons with Parkinson's disease and healthy elders. *Gait Posture*. 2004;19:215-225.

Dibble LE, Foreman KB, Addison O, et al. Exercise and medical effects on persons with Parkinson disease across the domains of disability: a randomized clinical trial. *J Neurol Phys Ther*. 2015;39:85-92.

Dichgans J, Diener HC. Different forms of postural ataxia in patients with cerebellar diseases. In: Igarashi M, Black FO, eds. *Disorders of Posture and Gait*. Amsterdam: Elsevier; 1986:207-213.

Dobkin B, Barbeau H, Basso M, et al. Weight-supported treadmill vs over-ground training for walking after acute incomplete SCI. *Neurol.* 2006;66:484-493.

Doorenbosch CAM, Harlaar J, Roebroek ME, Lankhorst GJ. Two strategies of transferring from sit-to-stand; the activation of monoarticular and biarticular muscles. *J Biomechanics.* 1994;27(11):1299-1307.

Dubost V, Beauchet O, Manckoundia P, et al. Decreased trunk angular displacement during sitting down: an early feature of aging. *Phys Ther.* 2005;85(5):404-412.

Duncan PW, Zorowitz R, Bates B, et al. Management of adult stroke rehabilitation care: a clinical practice guideline. *Stroke.* 2005;36:e100-e143.

Ebersbach G, Ebersbach A, Edler D, et al. Comparing exercise in Parkinson's disease – the Berlin LSVT® BIG study. *Movement Disorders.* 2010;25(12):1902-1908.

Ebersbach G, Grust U, Ebersbach A, et al. Amplitude-oriented exercise in Parkinson's disease: a randomized study comparing LSVT-BIG and short training protocol. *J Neural Transm.* 2015;122:253-256.

Ebersbach G, Ebersbach A, Gandor F, et al. Impact of physical exercise on reaction time in patients with Parkinson's disease—data from the Berlin BIG study. *Arch Phys Med Rehabil.* 2014;95:996-999.

Elstein AS. Thinking about diagnostic thinking: a 30-year perspective. *Adv in Health Sci Educ.* 2009;14:7-18.

Elstein AS and Schwarz A. Clinical problem solving and diagnostic decision making”: selective review of cognitive literature. *BMJ.* 2002;324:729-732.

Etnyre B and Thomas DQ. Event standardization of sit-to-stand movement. *Phys Ther.* 2007;87:1651-1666.

Farley BG and Koshland GF. Training BIG to move faster: the application of the speed-amplitude relation as a rehabilitation strategy for people with Parkinson's disease. *Exp Brain Res.* 20015;167(3):462-467.

Farley BG, Fox CM, Ramig LO, et al. Intensive amplitude-specific therapeutic approaches for Parkinsons disease: toward a neuroplasticity-principled rehabilitation model. *Topics in Geriatric Rehabil.* 2008;24(2):99-114.

Fasano A and Bloem BR. Gait disorders. *Movement Disorders.* 2013 Oct;19(5):1344-82.

Fernandez del Olmo M, Arias P, Furio MC, et al. Evaluation of the effect of training using auditory stimulation on rhythmic movement in Parkinsonian patients—a combined motor and [18F]-FDG PET study. *Parkinsonism Related Disorders.* 2006;12:155-164.

Freeland RL, Festa C, Sealy M, et al. The effects of pulsed auditory stimulation on various gait measurements in persons with Parkinson's disease. *NeuroRehabil.* 2002;17:81-87.

Garber CE, Blissmer B, Deschenes MR, et al. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exercise.* 2011;43(7):1334-59.

Giladi N, Horak FB, Hausdorff JM. Classification of gait disturbances: distinguishing between continuous and episodic changes. *Movement Disorders.* 2013;28(11):1469-1473.

Goldstein MS, Scalziti DA, Craik RL, et al. The revised research agenda for physical therapy. *Phys Ther.* 2011 Feb;91(2):165-74.

Graber ML. Educational strategies to reduce diagnostic error: can you teach this stuff? *Adv in Health Sci Educ.* 2009;14:63-69.

Guide to Physical Therapist Practice, 2<sup>nd</sup> ed. *Phys Ther.* 2001:81.

Guide to Physical Therapist Practice.  
[http://www.apta.org/uploadedFiles/APTAorg/About\\_Us/Policies/Practice/DocumentationPatientClientManagement](http://www.apta.org/uploadedFiles/APTAorg/About_Us/Policies/Practice/DocumentationPatientClientManagement)

.pdf. Accessed February, 2017.

Guccione AA. Physical therapy diagnosis and the relationship between impairments and function. *Phys Ther.* 1991;71(7):499-503.

Guccione, A.A., Wong, R.A., Avers, D. (2012). *Geriatric Physical Therapy*. St Louis, Missouri: Mosby, Inc.

Hackney ME and Earhart GM. Effects of dance on movement control in Parkinson's disease: a comparison of Argentine Tango and American ballroom. *J Rehabil Med.* 2009;41:475-481.

Harvey RL, Winstein CJ. Design for the Everest randomized trial of cortical stimulation and rehabilitation for arm function following stroke. *Neurorehabil Neural Repair.* 2009;23:32-44.

Hedman LD, Rogers MW, Hanke TA. Neurologic professional education: linking the foundation science of motor control with physical therapy interventions for movement dysfunction. *Neuro Rep* 1996;20(1):9-13.

Hedman LD, Morris DM, Graham CL, et al. Locomotor Requirements for Bipedal Locomotion: A Delphi Survey. *Phys. Ther.* 2014;94(1):52-67.

Hershberg JA, Rose DK, Tilson JL, et al. The interface of clinical decision-making with study protocols for knowledge translation from a walking recovery trial. *J Neurol Phys Ther.* 2017;41:59-67.

Hidler J, Nichols D, Pelliccio, et al. Multicenter randomized clinical trial evaluating the effectiveness of the Lokomat in subacute stroke. *Neurorehabil Neural Repair.* 2009.

Holleran CL, Straube DD, Kinnaird CR, et al. Feasibility and potential efficacy of high-intensity stepping training in variable context in subacute and chronic stroke. *Neurorehabil Neural Repair.* 2014;28(7):643-651.

Horak FB, Wrisley DM, Frank J. The balance evaluation systems test to differentiate balance deficits. *Phys Ther.* 2009;89(5):484-498.

Horn SD, DeJong G, Smout RJ, et al. Stroke rehabilitation patients, practice, and outcomes: is earlier and more aggressive therapy better? *Arch Phys Med Rehabil.* 2005;86(2):S101-14.

Hornby GT, Holleran CL, Hennessy PW, et al. Variable intensive early walking poststroke (VIEWS): a randomized controlled trial. *Neurorehabil Neural Repair.* 2016;30(5):440-450.

Hove MJ and Keller PE. Impaired movement timing in neurological disorders: rehabilitation and treatment strategies. *Ann N Y Acad Sci.* 2015;1337:111-117.

Janssen WGM, Bussman HBJ, Stam HJ. Determinants of the sit-to-stand movement: a review. *Phys Ther.* 2002;82:866-879.

Jette DU, Warren RL, Wirtalla C. The relation between therapy intensity and outcomes of rehabilitation in skilled nursing facilities. *Arch Phys Med Rehabil.* 2005;86(3):373-379.

Jones TE, Stephenson KW, King JG, et al. Sarcopenia – mechanisms and treatments. *J Geri Phys Ther.* 2009;32(9):39-45.

Kauffman, T.L., Barr, J.O., Moran, M.L. (2007). *Geriatric Rehabilitation Manual*. Philadelphia, PA: Elsevier

Kilmer DD. Response to resistive strengthening exercise training in humans with neuromuscular disease. *Am J Phys Med Rehabil.* 2002;81(11 Suppl):S121-S126.

Kimberley TJ, Samargia S, Moore LG, et al. Comparison of amounts and types of practice during rehabilitation for traumatic brain injury and stroke. *J Rehabil Res Dev.* 2010;47(9):851-62.

Kleim JA, Jones TA. Principles of experience-dependent neural plasticity: implications for rehabilitation after brain damage. *J Speech Lang Hearing Res.* 2008;51:s225-s239.

Kosak MC, Reding MJ. Comparison of partial body weight-supported treadmill gait training versus aggressive bracing assisted walking post stroke. *Neurorehabilitation and Neural Repair.* 2000;14:13-19.

Kwakkel G, Kollen BJ, Wagenaar RC. Long term effects of intensity of upper and lower limb training after stroke: a randomized trial. *J Neurol Neurosurg Psychiatry.* 2002;72:473-479.

Landi F, Marzetti Em, Martone AM, et al. Exercise as a remedy for sarcopenia. *Curr Opin Clin Nutr Metab Care.* 2014;17(1):25-31.

Lang CE and Bastian AJ. Additional somatosensory information does not improve cerebellar adaptation during catching. *Clin Neurophysiol.* 2001;112:895-907.

Lang CE, MacDonald JR, Reisman DS, et al. Observation of amounts of movement practice provided during stroke rehabilitation. *Arch Phys Med Rehabil.* 2009;90:1692-8.

Larsen DS. A career influenced by the STEP conferences. *J Neurol Phys Ther.* 2017;41:68-69.

Lee MJ, Kilbreath SL, Singh MF, et al. Effect of progressive resistance training on muscle performance after chronic stroke. *Med Sci Sports Exerc.* 2010;42(1):23-34.

Lim I, van Wegen E, deGoede C. Effects of external rhythmical cueing on gait in patients with Parkinson's disease: a systematic review. *Clin Rehabil.* 2005;19:695-713.

LSVT® BIG Training and Certification Workshop Binder. Copyright 2009 LSVT Global, LLC.

Macko RF, Ivey FM, Forrester LW. Task-oriented aerobic exercise in chronic hemiparetic stroke: training protocols and treatment effects. *Top Stroke Rehabil.* 2005;12:45-57.

Manckoundia P, Mourey F, Pfitzenmeyer P, et al. Is backward disequilibrium in the elderly caused by an abnormal perception of verticality? A pilot study. *Clin Neurophysiol.* 2007;118:786-793.

Manckoundia P, Mourey F, Perennou D, Pfitzenmeyer R. Backward disequilibrium in elderly subjects. *Clin Interventions Aging.* 2008;3(4):667-672.

Manckoundia P, Gerbault N, Mourey F, et al. Multidisciplinary management in geriatric day-hospital is beneficial for elderly fallers: a prospective study of 28 cases. *Arch Gerontol Geriatri.* 2007;44:61-70.

Malouin F, Ptvin M, Prevost J. et al. Use of an intensive task-oriented training program in a series of patients with acute cerebrovascular accidents. *Phys Ther.* 1992;72:781-794.

Marchese R, Diverio M, Zucchi F, et al. The role of sensory cues in the rehabilitation of Parkinsonian patients: a comparison of two physical therapy protocols. *Move Disorders.* 2000;15:879-883.

Martin K, Inman J, Kirschner A, et al. Characteristics of hypotonia in children: a consensus opinion of pediatric occupational and physical therapists. *Pediatr Phys Ther.* 2005;17:275-282.

Martone AM, Lattanzio F, Abbatecola AM, et al. Treating sarcopenia in older and oldest old. *Curr Pharm Des.* 2015;21(13):1715-1722.

Matheron E, Dubost V, Mourey F, et al. Analysis of postural control in elderly subjects suffering from Psychomotor Disadaptation Syndrome (PDS). *Arch Gerontol Geriatr.* 2010;51:e19-e23.

- Maughan KK, Lowry KA, Franke WD, Smiley-Oyen AL. The dose-response relationship of balance training in physically active older adults. *J Aging Phys Act.* 2012;20(4):442-55.
- Metter EL, Conwit R, Tobin J et al. Age-associated loss of power and strength in the upper extremities in women and men. *J Gerontol A Biol Sci Med Sci.* 1997;52:b267-B276.
- Millington PJ, Myklebust BM, Shambes GM. Biomechanical analysis of sit-to-stand motion in elderly persons. *Arch Phys Med Rehabil.* 1992;73:609-617.
- Montero-Fernandez N, Serra-Rexach JA. Role of exercise on sarcopenia in the elderly. *Eur J Phys Rehabil Med.* 2013;49(1):131-143.
- Morris ME. Locomotor training in people with Parkinson disease. *Phys Ther.* 2006;86:1426-1435.
- Morris ME. Movement disorders in people with Parkinson disease: a model for physical therapy. *Phys Ther.* 2000;80:578-597.
- Mourey F, Manckoundia P, Martin-Arveux I et al. Psychomotor disadaptation syndrome: a new clinical entity in geriatric patients. *Geriatrics.* 2004;59(5):20-24.
- Mourey F, Camus A, d'Athis P, et al. Mini motor test: a clinical test for rehabilitation of patients showing psychomotor disadaptation syndrome (PDS). *Arch Gerontol Geriatri.* 2005;40:201-211.
- Mudge S, Barber PA, Stott NS. Circuit-based rehabilitation improves gait endurance but not usual walking activity in chronic stroke: a randomized controlled trial. *Arch Phys Med Rehabil.* 2009;90(12):1989-96.
- Munin MC, Putman K, Hsieh CH, et al. Analysis of rehabilitation activities within skilled nursing and inpatient rehabilitation facilities after hip replacement for acute hip fracture. *Am J Phys Med Rehabil.* 2010;89(7):530-40.
- Nadeau SE, Wu SS, Dobkin BH, et al. Effects of task-specific and impairment-based training compared with usual care on functional walking ability after inpatient stroke rehabilitation: LEAPS trial. *Neurorehabil Neural Repair.* 2013;27(4):370-380.
- Nieuwboer A, Feys P, de Weerd W, et al. Is using a cue the clue to the treatment of freezing in Parkinson's disease? *Physiother Res Int.* 1997;2:125-134.
- Norton BJ. Clinical decision making in physical therapy: role of measurement and research in clinical decision making. In, *Saunders Manual of Physical Therapy Practice.* Saunders, 1995.
- Norton BJ. "Harnessing our collective professional power": diagnosis dialog. *Phys Ther.* 2007;87(6):635-638.
- Norton BJ. Diagnosis Dialog: progress report [guest editorial]. *Phys Ther.* 2007;87:1270-1273.
- Nuzik S, Lamb R, VanSant A, et al. Sit-to-stand movement pattern: a kinematic study. *Phys Ther.* 1986;66(11):1708-1713.
- Ouellette MM, LeBrasseur NK, Bean JF, et al. High-intensity resistance training improves muscle strength, self-reported function, and disability in long-term stroke survivors. *Stroke.* 2004;35:1404-1409.
- Pai Y-C and Rogers MW. Speed variation and resultant joint torques during sit-to-stand. *Arch Phys Med Rehabil.* 1991;72:881-885.
- Pak S, Patten C. Strengthening to promote functional recovery poststroke: an evidence-based review. *Topics Stroke Rehabil.* 2008;15(3):177-199.

- Patten C, Dozono J, et al. Combined functional task practice and dynamic high intensity resistance training promotes recovery of upper-extremity motor function in post-stroke hemiparesis: a case study. *J Neurol Phys Ther.* 30(3):99-119.
- Perennou DA, Mazibrada G, Chauvineau V, et al. Lateropulsion, pushing and verticality perception in hemisphere stroke: a causal relationship? *Brain.* 2008;131:2401-2413.
- Perennou DA, Amblard B, Leblond C, Pelissier J. Biased postural vertical in humans with hemispheric cerebral lesions. *Neurosci Letters.* 1998;252:75-78.
- Peterson MD, Rhea MR, Sen A et al. Resistance exercise for muscular strength in older adults: a meta analysis. *Aging Res Rev.* 2010;8(3):226-237.
- Pfitzenmeyer P, Mourey F, Tavernier B, Camus A. Psychomotor disadaptation syndrome. *Arch Gerontol Geriatr.* 1999;28:217-225.
- Pfitzenmeyer P, Mourey F, Mischis-Troussard C, Bonneval P. Rehabilitation of serious postural insufficiency after falling in very elderly subjects. *Arch Gerontol Geriatr.* 2001;33:211-218.
- Pfitzenmeyer P, Mourey F, Manckoundia P, d'Athis P. A 4-year follow-up of very old patients presenting with frontal-subcortical dysfunction compared with Alzheimer's disease patient. *Gerontol.* 2005;51:62-65.
- Phu S, Boersma D, Duque G. Exercise and Sarcopenia. *J Clin Densitom.* 2015. 18(4):488-492.
- Physical Therapist Practice and The Human Movement System White Paper, August 2015, American Physical Therapy Association.
- Porter M and Lee T. The Strategy That Will Fix Health Care. *Harvard Business Review.* 2013:October.
- Protas EJ, Mitchell K, Williams A, et al. Gait and step training to reduce falls in Parkinson's disease. *NeuroRehabil.* 2005;20:183-190.
- Pugh KG and Lipsitz LA. The microvascular frontal-subcortical syndrome of aging. *Neurobio Aging.* 2002;23:421-431.
- Quinn & Gordon (2016). *Documentation for Rehabilitation: A guide to clinical decision making in rehabilitation.* 3rd Ed. Elsevier: Philadelphia.
- Quinn L and Busse M. Physiotherapy clinical guidelines for Huntington's disease. *Neurodegen Dis Manage.* 2012;2(1):1-11.
- Raymond MJ, Bramley-Tzerfos RE, Jeffs KJ, et al. Systematic review of high-intensity progressive resistance strength training of the lower limb compared with other intensities of strength training in older adults. *Arch Phys Med Rehabil.* 2013;94(8):1458-1472.
- Richards CL, Malouin F, Wood-Dauphinee S, et al. Task-specific physical therapy for optimization of gait recover in acute stroke patients. *Arch Phys Med Rehabil.* 1993;74:612-620.
- Roerdink M, Lamoth CJC, Kwakkel G, et al. Gait coordination after stroke: benefits of acoustically paced treadmill walking. *Phys Ther.* 2007;87:1009-1022.
- Romero-Arenas S, Martinez-Pascual M, Alcaraz PE. Impact of resistance circuit training on neuromuscular, cardiorespiratory and body composition adaptations in the elderly. *Aging Dis.* 2013;4(5):256-263.
- Rose SJ. Physical therapy diagnosis: role and function. *Phys Ther.* 1989;69:535-537.
- Rothstein JM, Echternach JL, Riddle DL. The hypothesis-oriented algorithm for clinicians II (HOACII): a guide for patient management. *Phys Ther.* 2003;83(5):455-470.

Rubinstein TC, Giladi N, and Hausdorff JM. The power of cueing to circumvent Dopamine deficits: a review of physical therapy treatment of gait disturbances in Parkinson's disease. *Movement Disorders*. 2002;17:1148-1160.

Sahrmann SA. Diagnosis by the physical therapist-a prerequisite for treatment. A special communication. *Phys Ther*. 1988;68(11):1703-1706.

Sahrmann SA. The human movement system: our professional identity. *Phys Ther*. 2014;94(7):1034-1042.

Sahrmann SA. The twenty-ninth Mary McMillan lecture: Moving precisely? Or taking the path of least resistance? *Phys Ther*. 1998;78(11):1208-1218.

Salbach NM, Mayo NE, Wood-Dauphinee S. et al. A task-oriented intervention enhances walking distance and speed in the first year post stroke: a randomized controlled trial. *Clin Rehabil*. 2004;18:509-519.

Schaefer RS. Auditory rhythmic cueing in movement rehabilitation: findings and possible mechanisms. *Phil Trans R Soc B*. 369: 20130402. <http://dx.doi.org/10.1098/rstb.2013.0402>.

Schaefer SY, Patterson CB, Lang CE. Transfer of training between distinct motor tasks after stroke: implications for task-specific approaches to upper-extremity neurorehabilitation. *Neurorehabil Neural Repair*. 2013 Apr 2 [Epub ahead of print].

Schauer M and Mauritz K-H. Musical motor feedback (MMF) in walking hemiparetic stroke patients: randomized trials of gait improvement. *Clin Rehabil*. 2003;17:713-722.

Scheets, Sahrmann, Norton. *Neurol Report*. 1999;23:158-169.

Scheets PL, Sahrmann SA, Norton BJ. Use of movement system diagnoses in the management of patients with neuromuscular conditions: a multiple-patient case report. *Phys Ther*. 2007;87(6):654-669.

Scheets PL, Sahrmann SA, Norton BJ, Stith JS, Crouner BE. What is backward disequilibrium and how do I treat it: a complex patient case study. *J Neurol Phys Ther*. 2015;39:119-126.

Schindl MR, Forstner C, Kern H, Hesse S. Treadmill training with partial body weight support in nonambulatory patients with cerebral palsy. *Arch Phys Med Rehabil*. 2000;81:301-306.

Schultz AB, Alexander NB, Ashton-Miller JA. Biomechanical analyses of rising from a chair. *J Biomechanics*. 1992;25(12):1383-1391.

Silva NL, Oliveira RB, FleckSJ, et al. Influence of strength training variables on strength gains in adults over 55 years-old: a meta-analysis of dose-response relationships. *J Sci Med Sport*. 2014;17(3):337-344.

Sousa N, Mendes R, Monteiro G, et al. Progressive resistance strength training and the related injuries in older adults: the susceptibility of the shoulder. *Aging Clin Exp Res*. 2014;26(3):235-240.

Staab JP. Chronic subjective dizziness continuum. *Neurootology*. 2012;18(5):1118-11141.

Sullivan KJ, Brown DA, Klassen T, et al. Effects of task-specific locomotor and strength training in adults who were ambulatory after stroke: results of the STEPS randomized clinical trial. *Phys Ther*. 2007;87:1580-1602.

Suteerawattananon M, Morris GS, Etnyre BR, et al. Effects of visual and auditory cues on gait in individuals with Parkinson's disease. *J Neurological Sci*. 2004;219:63-69.

Taylor NF, Dodd KJ, Damiano DL. Progressive resistance exercise in physical therapy: a summary of systematic reviews. *Phys Ther*. 2005;85(11):1208-1223.

Thaut MH, Leins AK, Rice RR, et al. Rhythmic auditory stimulation improves gait more than NDT/Bobath training in near-ambulatory patients early poststroke: a single-blind randomized trial. *Neurorehabil Neural Repair*. 2007;21:455-459.

Tilson JK, Settle SM, Sullivan KJ. Application of evidence-based practice strategies: current trends in walking recovery interventions poststroke. *Topics Stroke Rehabil*. 2008;15(3):227-246.

Valenzuela T. Efficacy of progressive resistance training interventions in older adults in nursing homes: a systematic review. *J Am Med Dir Assoc*. 2012;13(5):418-28.

Wennberg J, Gittelsohn A. Small area variations in healthcare delivery. *Science*. 1973;182(117):1102-8.

Williams G, Lai D, Schache A, Morris ME. Classification of gait disorders following traumatic brain injury. *J Head Trauma Rehabil*. 2015;30(2):E13-E23.

Wittwer JE, Webster KE, Hill K. Rhythmic auditory cueing to improve walking in patients with neurological conditions other than Parkinson's disease – what is the evidence? *Disabil Rehabil*. 2013;35(2):164-176.

Wrentenberg P, Lindberg F, Arborelius UP. Effect of armrests and different ways of using them on hip and knee load during rising. *Clin Biomech*. 1993;8:95-101.