

# SCHIZOPHRENIA IN LONG-TERM CARE



Douglas Steenblock, MD

Iowa Veterans Home



# DISCLOSURES



➤ NONE

# OBJECTIVES



- Discuss the psychopharmacologic management of schizophrenic patients in long-term care settings.
- Identify the unique clinical aspects of schizophrenia in late life.

# SCHIZOPHRENIA



- “*Dementia Praecox*”
- Lifetime prevalence of 1% (male=female)
  - 0.1-0.5% of elderly
- Peak age of onset:
  - 10-25 (men)
  - 25-35 (women)
- Ventricular enlargement, cortical atrophy and soft neurological signs.
- *Prodrome* often occurs prior to onset.





# AGE OF ONSET

- Early onset: Up to age 40 (45)
  - 75-80% of cases
  
- Late onset: 40-60
  - Often female
  - Better prognosis
  
- Very late onset: 60+
  - Rare
  - More medical co-morbidities

# SCHIZOPHRENIA



- Cause unknown.
  
- Risk factors:
  - Genetics.
    - 50% concordance monozygotic twins.
  - Winter/spring birth.
  - Influenza during pregnancy.
  
- Dopamine Hypothesis:
  - Increased limbic, decreased frontal.
  - Other neurotransmitters also implicated

# SCHIZOPHRENIA: DIAGNOSTIC CRITERIA



- 2 or more of following actively occurring over 1 month period:
  - Delusions\*
  - Hallucinations\*
  - Disorganized speech\*
  - Grossly disorganized or catatonic behavior
  - Negative symptoms
  
- Total duration of illness 6 months or more

# SCHIZOPHRENIA SYMPTOMS



## ➤ Positive Symptoms:

- Delusions
- Hallucinations
- Disorganized speech
- Disorganized behavior

## ➤ Negative Symptoms:

- Affect flat
- Alogia (mute)
- Avolition
- Autistic
- Apathy
- Ambivalence
- Anhedonia
- Etc.



# PSYCHOTIC SYMPTOMS



## Schizophrenia:

- Bizarre or complex delusions
- Auditory hallucinations
- Long-term

## Dementia-related psychosis:

- Misidentification
- Well-formed visual hallucinations
- Usually remits

# COGNITIVE IMPAIRMENT



- More than 70% of schizophrenics affected.
- May be less impactful for late-onset cases.
- May be worse in institutionalized vs. community.
- May affect executive function, memory, IQ, visuospatial function, processing speed, attention, arithmetic and verbal fluency.
- Cognitive abilities may remain stable over the course of the illness in many patients.
  - Those in nursing homes more likely to decline.
- Schizophrenics overall have higher risk of dementia.

# SCHIZOPHRENIA TREATMENT



## ➤ Antipsychotic medication:

- Conventional/Typical (1<sup>st</sup> Gen)
  - Dopamine (D2) antagonist
- Novel/Atypical (2<sup>nd</sup> Gen)
  - Dopamine & serotonin antagonist

## ➤ ECT

- Safe and effective option for older schizophrenics.

## ➤ Psychosocial:

- Social skills training, supported employment, cognitive remediation, psychotherapy, groups.
- Family support and case management.



# ANTIPSYCHOTICS

## First Gen (Typical)

- Haloperidol (Haldol)
- Fluphenazine (Prolixin)
- Thiothixine (Navane)
- Perphenazine (Trilafon)
- Trifluoperzaine (Stelazine)
- Thioridazine (Mellaril)
- Chlorpromazine (Thorazine)
- Loxapine (Loxitane)
- Others

## Second Gen (Atypical)

- Risperidone (Risperdal)\*
- Olanzapine (Zyprexa)\*
- Clozapine (Clozaril)
- Quetiapine(Seroquel)
- Ziprasidone (Geodon)
- Aripiprazole (Abilify)
- Paliperidone (Invega)
- Lurasidone (Latuda)
- Asenapine (Saphris)
- Iloperidone (Fanapt)
- Others

\* (Trade names in parentheses)

# EXTRAPYRAMIDAL SIDE EFFECTS (EPS)



## Parkinsonism:

- 20-35% overall prevalence
  - Higher for elderly
- Resting tremor
- Rigidity
- Bradykinesia
- Shuffling gait
  
- Treat with anticholinergic or amantadine.

## Other:

- Dystonia (torticollis, oculogyric)
- Drooling
- Akathisia (restlessness)
- Tardive dyskinesia



# TARDIVE DYKINESIA

- Abnormal movements of face/mouth such as lip smacking, clenching, chewing, puckering, and tongue thrusting. May also see choreiform movements in limbs.
- Prevalence 20-30% overall (50-60% for elderly).
- Usually develops over months or years, but sometimes sooner.
- Most cases are irreversible (dopamine receptor sensitivity altered).
- Risk factors: Female, higher negative/positive symptoms, greater cognitive impairment, other drug-induced movements, substance use, diabetes.

# TARDIVE DYSKINESIA: MANAGEMENT



- Prevention: Screening is crucial.
- Choose antipsychotics with less risk. Once TD starts, switching may be ineffective.
- Anticholinergics may worsen; try to eliminate.
- Reduction in dose may worsen.
- Consider severity and impact on patient.
- VAMT2 inhibitors:
  - Valbenazine
  - Deutetrabenazine



# RELATIVE RISK OF EPS FOR ATYPICAL ANTIPSYCHOTICS

<u>Risk:</u>	<u>Agents:</u>
Low	Clozapine Quetiapine
Intermediate	Olanzapine
High	Risperidone Ziprasidone Aripiprazole



# NEUROLEPTIC MALIGNANT SYNDROME (NMS)



- Associated with initiation of or increase in antipsychotic.
- High doses or multiple antipsychotics have higher risk.
- **Features:** Confusion, delirium, tremor, stiffness, autonomic instability, fever, death.
- **Creatinine Kinase** markedly elevated; WBC and LFTs may be elevated as well.
- **Tx:** Stop antipsychotic, admit to medical, supportive measures, may need bromocriptine , dantrolene or amantadine.



# ANTICHOLINERGIC SIDE EFFECTS

- Increased confusion
- Dry mouth
- Blurred vision
- Constipation
- Urine retention



# METABOLIC SYNDROME

- Mainly associated with second generation (atypical) agents.
- Weight gain, hyperlipidemia, hyperglycemia, higher risk of diabetes.
- 32.5% of schizophrenics.
- Metformin may be beneficial.
- **Risk for weight gain:**
  - **Higher Risk:**
    - Olanzapine, clozapine
  - **Medium Risk:**
    - Risperidone, quetiapine, paliperidone
  - **Lower Risk:**
    - Aripiprazole, ziprasidone, lurasidone, asenapine, haloperidol

# OTHER COMPLICATIONS



- Orthostatic hypotension (falls)
- Hyperprolactinemia
- Cardiac conduction (QT prolongation)
- Seizures
- Antihistaminic:
  - Sedation
  - Weight gain

# CLOZAPINE



- May be used in cases of EPS sensitivity or non-response to treatment.
  
- Requires monitoring of WBC/ANC due to risk of **agranulocytosis** (q 1-4 weeks).
  
- Other possible liabilities:
  - Seizure risk
  - Metabolic
  - Cardiomyopathy
  - Anticholinergic

# LONG-TERM INJECTABLE ANTIPSYCHOTICS



- Haloperidol, risperidone, paliperidone, aripiprazole and others.
- Intervals vary from 2 weeks to 3 months.
- Outcomes in controlled settings may not be superior to oral.

# DEPRESSION



- May be difficult to differentiate from negative symptoms.
- Rule out medical conditions that may mimic depression.
- Not particular agent favored.
- Use atypical antipsychotics.
- Consider psychotherapy.
- Important to treat:
  - Cognition
  - Suicide risk

# SUICIDE RISK



- Suicidal ideation prevalence: 10% current, 56% lifetime.
- Risk factors: Multiple hospitalizations, previous attempts, co-morbid mood and personality disorders, substance abuse.
- Major depression + schizophrenia = greatly increased risk



# SCHIZOPHRENIA: PROGNOSIS



- About 60% of patients have substantial improvement or full remission of positive symptoms on antipsychotic medication.
- Many experience slow functional deterioration over lifetime.
  - Most will not be able to work (unless supported).
  - Less likely to drive, marry, or live independently than bipolar cohorts.
- Lower socioeconomic status: “*Downward Drift*”

# SCHIZOPHRENIA-SPECTRUM DISORDERS



# SCHIZOPHRENIFORM DISORDER



- Same criteria for Schizophrenia, but total duration is less than 6 months.
- Often rapid onset.
- Prognosis is better.
- Treated with antipsychotics, but may not need long term.

# SCHIZOAFFECTIVE DISORDER



- Elements of schizophrenia combined with elements of either bipolar disorder or major depression.
  
- The mood Sx (mania and/or depression) must account for a substantial portion of the total duration of the illness.
  
- Patient has periods (2 weeks or more) of psychosis occurring during times when they are not manic or depressed.
  
- Bipolar/Depressive Types.
  - With catatonia



# SCHIZOAFFECTIVE DISORDER: TREATMENT

- Similar to schizophrenia, but also mood stabilizers and antidepressants.
- Prognosis similar to schizophrenia

# SCHIZOTYPAL PERSONALITY DISORDER



- A pervasive pattern of social and interpersonal deficits marked by acute discomfort with, and reduced capacity for, close relationships as well as by cognitive or perceptual distortions and eccentricities of behavior, beginning by early adulthood and present in a variety of contexts, as indicated by five (or more) of the following:
  - Ideas of reference (excluding delusions).
  - Odd beliefs or magical thinking that influences behavior and is inconsistent with subcultural norms.
  - Unusual perceptual experiences, including bodily illusions.
  - Odd thinking and speech.
  - Suspiciousness or paranoid ideation.
  - Inappropriate or constricted affect.
  - Behavior or appearance that is odd, eccentric, or peculiar.
  - Lack of close friends or confidants other than first-degree relatives.
  - Excessive social anxiety that does not diminish with familiarity and tends to be associated with paranoid fears rather than negative judgments about self.
  
- Late-onset schizophrenics are often diagnosed with cluster A personality disorders prior to the onset of schizophrenia.

# ISSUES IN LONG-TERM CARE



# NURSING HOMES



- Have become defacto institutions for those with serious mental illness (SMI), following deinstitutionalization and closing of state hospitals.
- In VA system, as inpatient beds and stays decreased, prevalence of SMI in nursing homes increased.
- 2.7-7.1% of nursing home residents have SMI.
- >80% of schizophrenics live in the community.



# NURSING HOMES



- **Residents with SMI more likely to stay long term.**
- **Residents with SMI are younger:**
  - Average age at time of entry is 65 for SMI, and 80 for those who are not SMI.
  - Middle aged patients with schizophrenia are 4 times more likely to need nursing home placement than same-age peers without SMI.
- **Nursing homes with larger percentage of SMI residents tend to be lower in quality:**
  - Staffed at lower levels, higher Medicaid, use restraints more, more ulcers, more feeding tubes, more catheters, more hospitalizations, more deficiency citations.

# NURSING HOMES



- Compared with community cohorts, patients with SMI in nursing homes are more likely to have severe psychiatric symptoms, greater cognitive deficits, more functional and physical impairment, more aggressive behavior, and less social support.
- In nursing homes, those with dementia alone had more aggressive behavior than those who had SMI alone. But those who had dementia AND SMI had the highest rates of all.
- **Challenges:** Training for staff, finding psychiatric providers, developing appropriate activities for younger residents.

# CO-MORBID MEDICAL CONDITIONS



- Tend to be exacerbated by schizophrenia.
- Schizophrenics often receive inadequate healthcare.
- Medications may have long-term effects.
- Substance use (including tobacco) may contribute.

# END-OF-LIFE CARE FOR SCHIZOPHRENICS



- More likely than cohorts to die in nursing homes.
- Less likely to see (non-psychiatric) specialists.
- Less likely to be prescribed analgesics.
- Less likely to receive palliative care.
- Psychiatric medications should be continued as comfort care.



# ANTIPSYCHOTIC REDUCTION

- Length of treatment guidelines vary for all age groups.
- Very little research on reductions for older schizophrenics.
- Risk/benefit analysis is needed (including patient preference).
- Patients on the ends of the severity spectrum may be best candidates.



# ANTIPSYCHOTIC REDUCTION

- For all age groups, following factors are associated with lower rate of relapse after discontinuation:
  - Low dose prior to discontinuation.
  - Older age.
  - Shorter duration of untreated psychosis.
  - Lower severity of positive symptoms.
  - Better social functioning.
  - Lower number of previous relapses.
  - Later onset.

# QUESTIONS?



## References:

1. Amad A, Geoffroy PA, Vaiva G, Thomas P. [Personality and personality disorders in the elderly: diagnostic, course and management]. *Encephale*. 2013 Oct;39(5):374-82. doi: 10.1016/j.encep.2012.08.006. Epub 2012 Oct 24. Review.
2. Andrews AO, Bartels SJ, Xie H, Peacock WJ. Increased risk of nursing home admission among middle aged and older adults with schizophrenia. *Am J Geriatr Psychiatry*. 2009 Aug;17(8):697-705..
3. Aschbrenner K, Grabowski DC, Cai S, Bartels SJ, Mor V. Nursing home admissions and long-stay conversions among persons with and without serious mental illness. *J Aging Soc Policy*. 2011 Jul-Sep;23(3):286-304.
4. Bartels SJ, Pratt SI. Psychosocial rehabilitation and quality of life for older adults with serious mental illness: recent findings and future research directions. *Curr Opin Psychiatry*. 2009 Jul;22(4):381-5.
5. Becker M, Andel R, Boaz T, Howell T. The association of individual and facility characteristics with psychiatric hospitalization among nursing home residents. *Int J Geriatr Psychiatry*. 2009 Mar;24(3):261-8.
6. Bowersox NW, Szymanski BJ, McCarthy JF. Associations between psychiatric inpatient bed supply and the prevalence of serious mental illness in Veterans Affairs nursing homes. *Am J Public Health*. 2013 Jul;103(7):1325-31.
7. Caroff SN. Overcoming barriers to effective management of tardive dyskinesia. *Neuropsychiatr Dis Treat*. 2019 Apr 4;15:785-794. doi: 10.2147/NDT.S196541. eCollection 2019. Review.
8. Cen X, Li Y, Hasselberg M, Caprio T, Conwell Y, Temkin-Greener H. Aggressive Behaviors Among Nursing Home Residents: Association With Dementia and Behavioral Health Disorders. *J Am Med Dir Assoc*. 2018 Dec;19(12):1104-1109.e4.
9. Chochinov HM, Martens PJ, Prior HJ, Kredentser MS. Comparative health care use patterns of people with schizophrenia near the end of life: a population-based study in Manitoba, Canada. *Schizophr Res*. 2012 Nov;141(2-3):241-6.
10. Citrome L. Clinical management of tardive dyskinesia: Five steps to success. *J Neurol Sci*. 2017 Dec 15;383:199-204. Nov 16. Review.
11. Dickerson FB. Women, aging, and schizophrenia. *J Women Aging*. 2007;19(1-2):49-61.
12. Essali A, Ali G. Antipsychotic drug treatment for elderly people with late-onset schizophrenia. *Cochrane Database Syst Rev*. 2012 Feb 15;(2):CD004162.
13. Felmet K, Zisook S, Kasckow JW. Elderly patients with schizophrenia and depression: diagnosis and treatment. *Clin Schizophr Relat Psychoses*. 2011 Jan;4(4):239-50.
14. Folsom DP, Lebowitz BD, Lindamer LA, Palmer BW, Patterson TL, Jeste DV. Schizophrenia in late life: emerging issues. *Dialogues Clin Neurosci*. 2006;8(1):45-52.
15. Mental Illness In Nursing Homes: Variations Across States. Grabowski DC, Aschbrenner KA, Feng Z, Mor V. *Health affairs (Project Hope)*. 2009 May-Jun; 28(3): 689-700.
16. Guo JY, Ragland JD, Carter CS. Memory and cognition in schizophrenia. *Mol Psychiatry*. 2019 May;24(5):633-642. 21.
17. Hsu HF, Kao CC, Lu T, Ying JC, Lee SY. Differences in the Effectiveness of Long-Acting Injection and Orally Administered Antipsychotics in Reducing Rehospitalization among Patients with Schizophrenia Receiving Home Care Services. *J Clin Med*. 2019 Jun 9;8(6). pii: E823.
18. Jalenques I, Rachez C, Tourtauchaux R, Cellier Y, Legrand G. [Old patients suffering from long-standing schizophrenia: clinical aspects]. *Geriatr Psychol Neuropsychiatr Vieil*. 2011 Sep;9(3):345-53. doi: 10.1684/pnv.2011.0285. Review.
19. Jester DJ, Hyer K, Bowblis JR. Quality Concerns in Nursing Homes That Serve Large Proportions of Residents With Serious Mental Illness. *Gerontologist*. 2020 Sep 15;60(7):1312-1321. 20. Karim S, Overshott R, Burns A. Older people with chronic schizophrenia. *Aging Ment Health*. 2005 Jul;9(4):315-24. Review. PubMed [citation] PMID: 16019287
21. Kasckow JW, Zisook S. Co-occurring depressive symptoms in the older patient with schizophrenia. *Drugs Aging*. 2008;25(8):631-47.
22. Kiosses DN, Szanto K, Alexopoulos GS. Suicide in older adults: the role of emotions and cognition. *Curr Psychiatry Rep*. 2014 Nov;16(11):495.
23. Kumagaya D, Halliday G. Acute electroconvulsive therapy in the elderly with schizophrenia and schizoaffective disorder: a literature review. *Australas Psychiatry*. 2019 Oct;27(5):472-476.



24. Kurtz MM. Neurocognitive impairment across the lifespan in schizophrenia: an update. *Schizophr Res.* 2005 Apr 1;74(1):15-26.
25. Li Y, Cai X, Cram P. Are patients with serious mental illness more likely to be admitted to nursing homes with more deficiencies in care? *Med Care.* 2011 Apr;49(4):397-405. doi: 10.1097/MLR.0b013e318202ac10.
26. López-Álvarez J, Sevilla-Llewellyn-Jones J, Agüera-Ortiz L. Anticholinergic Drugs in Geriatric Psychopharmacology. *Front Neurosci.* 2019 Dec 6;13:1309.
27. Maglione JE, Thomas SE, Jeste DV. Late-onset schizophrenia: do recent studies support categorizing LOS as a subtype of schizophrenia? *Curr Opin Psychiatry.* 2014 May;27(3):173-8.
28. Matei VP, Mihailescu AI, Davidson M. Is non-pharmacological treatment an option for certain schizophrenia patients? *Psychiatr Danub.* 2014 Dec;26(4):308-13.
29. McGarry BE, Joyce NR, McGuire TG, Mitchell SL, Bartels SJ, Grabowski DC. Association between High Proportions of Seriously Mentally Ill Nursing Home Residents and the Quality of Resident Care. *J Am Geriatr Soc.* 2019 Nov;67(11):2346-2352.
30. Mishara AL, Goldberg TE. A meta-analysis and critical review of the effects of conventional neuroleptic treatment on cognition in schizophrenia: opening a closed book. *Biol Psychiatry.* 2004 May 15;55(10):1013-22.
31. Muralidharan A, Mills WL, Evans DR, Fujii D, Molinari V. Preparing Long-Term Care Staff to Meet the Needs of Aging Persons With Serious Mental Illness. *J Am Med Dir Assoc.* 2019 Jun;20(6):683-688.
32. Murante T, Cohen CI. Cognitive Functioning in Older Adults With Schizophrenia. *Focus (Am Psychiatr Publ).* 2017 Jan;15(1):26-34.
33. Rahman M, Grabowski DC, Intrator O, Cai S, Mor V. Serious mental illness and nursing home quality of care. *Health Serv Res.* 2013 Aug;48(4):1279-98.
34. Rajji TK, Ismail Z, Mulsant BH. Age at onset and cognition in schizophrenia: meta-analysis. *Br J Psychiatry.* 2009 Oct;195(4):286-93.
35. Rajji TK, Miranda D, Mulsant BH. Cognition, function, and disability in patients with schizophrenia: a review of longitudinal studies. *Can J Psychiatry.* 2014 Jan;59(1):13-7.
36. Rajji TK, Mulsant BH. Nature and course of cognitive function in late-life schizophrenia: a systematic review. *Schizophr Res.* 2008 Jul;102(1-3):122-40.
37. Sfera A. Can neurostimulation prevent the risk of Alzheimer's disease in elderly individuals with schizophrenia? *Front Psychiatry.* 2013 Sep 18;4:111.
38. Shah JN, Qureshi SU, Jawaid A, Schulz PE. Is there evidence for late cognitive decline in chronic schizophrenia? *Psychiatr Q.* 2012 Jun;83(2):127-44.
39. Stegmayer K, Walther S, van Harten P. Tardive Dyskinesia Associated with Atypical Antipsychotics: Prevalence, Mechanisms and Management Strategies. *CNS Drugs.* 2018 Feb;32(2):135-147.
40. Suzuki T, Uchida H. Successful withdrawal from antipsychotic treatment in elderly male inpatients with schizophrenia—description of four cases and review of the literature. *Psychiatry Res.* 2014 Dec 15;220(1-2):152-7.
41. Tampi RR, Young J, Hoq R, Resnick K, Tampi DJ. Psychotic disorders in late life: a narrative review. *Ther Adv Psychopharmacol.* 2019 Oct 16;9:2045125319882798.
42. Tani H, Suzuki T, Wolfgang Fleischhacker W, Tomita M, Mimura M, Uchida H. Clinical Characteristics of Patients With Schizophrenia Who Successfully Discontinued Antipsychotics: A Literature Review. *J Clin Psychopharmacol.* 2018 Dec;38(6):582-589.
43. Temkin-Greener H, Campbell L, Cai X, Hasselberg MJ, Li Y. Are Post-Acute Patients with Behavioral Health Disorders Admitted to Lower-Quality Nursing Homes? *Am J Geriatr Psychiatry.* 2018 Jun;26(6):643-654.
44. Veerman SRT, Schulte PFJ, de Haan L. Treatment for Negative Symptoms in Schizophrenia: A Comprehensive Review. *Drugs.* 2017 Sep;77(13):1423-1459.
45. Ward KM, Citrome L. Antipsychotic-Related Movement Disorders: Drug-Induced Parkinsonism vs. Tardive Dyskinesia-Key Differences in Pathophysiology and Clinical Management. *Neurol Ther.* 2018 Dec;7(2):233-248.
46. Kaplan and Saddock's Synopsis of Psychiatry, 11th Edition, 2015