

Geriatric Behavioral Health Conference

Substance Use Disorders and the Brain In the Geriatric Population

Michelle Davids DO
Psychiatrist, Broadlawns Medical Center
ABPN Board Certified in Psychiatry and Addiction Psychiatry

Dhrumil Patel MD
Psychiatry Resident
Broadlawns Medical Center Unity Point Residency Program

1

Geriatric Behavioral Health Conference

DISCLOSURE

- Dr. Davids and Dr. Patel do not have any financial relationships with commercial interest companies to disclose.
- We will not be discussing off-label use of a commercial product.

2

Geriatric Behavioral Health Conference

Learning Objectives

- Define Substance Use Disorders
- Understand ways in which to screen substances use disorders in the geriatric population
- Common substances of misuse
- Treatment of substance use disorders in the geriatric population

3

Question

What is considered the size of one standard drink?

- A. 16 oz of regular beer
- B 8 oz of wine
- C 1.5 oz of distilled spirits
- D All of the above

4

What is a Substance Use Disorder?

- Per the DSM-5:
- Substance is often taken in larger amounts or over a longer period than intended
- Persistent desire or unsuccessful efforts to cut down or control use
- A great deal of time is spent in activities necessary to obtain the substance or recover from its effects

• American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Arlington, VA: American Psychiatric Publishing.

5

DSM-5 Continued

- Craving or a strong desire or urge to use
- Recurrent use resulting in a failure to fulfill major obligations at work, home, or school
- Continued use despite persistent or recurrent social or interpersonal problems caused or exacerbated by effects of use
- Important social, occupational, or recreational activities given up or reduced due to use

• American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Arlington, VA: American Psychiatric Publishing.

6

DSM-5 Continued

- Recurrent use in situations in which it is physically hazardous
- Use is continued despite knowledge of having ongoing or recurrent physical or psychological problems that are likely caused by or worsened by the substance use

7

DSM-5 Continued

- Tolerance: A need to use increased amounts of alcohol needed to achieve intoxication or desired effect OR a diminished effect with continued use of the same amount of a substance
- Withdrawal: varies based on substance. For alcohol: autonomic changes, tremor, insomnia, GI upset, hallucinations, agitation, anxiety, possible seizures

8

Question

Diagnosis of substance use disorder is divided in to 4 criteria. What criteria are needed?

- A. Impaired control
- B. Social Impairment
- C. Risky use
- D. Pharmacological Criteria
- E. All of the above

9

DSM-5 Continued

- Mild: 2-3 symptoms
- Moderate: 4-5 symptoms
- Severe: 6 or more symptoms

10

Substance Use Disorders in the Geriatric Population are often overlooked

- Patients are stereotyped as young
- Providers may be embarrassed to ask
- Patients may fear judgment and under report their use
- The baby boom generation is unique in its exposure to, attitudes toward, and prevalence of substance use. We should expect the rates of substance use to increase over the next twenty years.

Source: Kuerbis A, Sacco P, Blazer DG, Moore AA. Substance abuse among older adults. Clin Geriatr Med. 2014 Aug;30(3):629-54. doi: 10.1016/j.cger.2014.04.008. Epub 2014 Jun 12. PMID: 25037298; PMCID: PMC4146436.

11

Waves of Change

- In1900, individuals aged 65 years and older made up 1% of the global population, that proportion is estimated to top 20% in 2050

12

Why should we be concerned?

- Ongoing, undiagnosed substance use further complicates co-occurring medical problems
- Patients are at higher risk for falls and delirium
- Substance use worsens co-occurring psychiatric diagnosis and may increase the risk of suicide
- Older adults take more prescribed and over-the-counter medications than younger adults, increasing the risk for harmful drug interactions and misuse

Kennedy GJ, Ehrenová I, Frazer A, et al. The emerging problems of alcohol and substance abuse in late life. J Soc Distress Homeol. 1999;8(4):227-239

13

Standard Drink- World Health Organization

	Adult males <65 years old	Adult females (all ages) and adult males ≥65 years old
Abstinence	0 g	0 g
Low risk	1–40 g	1–20 g
Medium risk	41–60 g	21–40 g
High risk	61–100 g	41–60 g
Very high risk	101+ g	61+ g

The number of grams in a standard drink varies across countries. The US has one of the largest standard drinks (14 g), topped only by Austria (20 g), whereas Iceland and the UK have the smallest, with 8 g of ethanol considered to be a standard drink. The vast majority of countries across the world considers a standard drink to contain 10–12 g ethanol.

14

Substance Use Breakdown

- Out of all geriatric psychiatric patients with Substance Use Disorders admitted between 1999-2009:
 - 73.3% alcohol related disorders
 - 11% sedative-hypnotic use disorders
 - 2.9% opioid use disorders
 - 1% cannabis use disorders

Source: Dombrowski D, Norrell N, Holdroyd S. Substance use disorders in elderly admissions to an academic psychiatric inpatient service over a 10-year period. Journal of Addiction. Volume 2016, Article ID 4973018

15

Question

Which parts of the brain plays very important role in reward Circuit?

- A. Prefrontal cortex and Orbitofrontal cortex
- B. Anterior cingulate gyrus
- C. Raphe Nucleus
- D. Nucleus Accumbens and VTA
- E. Putamen and thalamus

16

Question

Which is the final common pathway for substance use disorder/addiction?

- A. MesoCortico-Limbic dopamine pathway
- B. Mesocortico pathway
- C. NigroStriatal pathway
- D. Tuberoinfundabular pathway
- E. Mesolimbic pathway

17

Neuroscience of how drugs hook people.

- Drug of abuse activates brain reward system.
- Reward—is defined as any event that increases the probability of response and has a positive hedonic effect.
- Principal focus of neurobiology of reporting effects of drugs with addiction potential has been the origins and terminal areas of mesocortico-limbic dopamine system

18

Neuroscience cont.

Neurotransmitter	Receptors	Effects	Exogenous
Glutamate	AMPA, NMDA	Excitatory	None
GABA	GABA _A , GABA _B	Inhibitory	None
Dopamine	D ₁ , D ₂	Excitatory/Inhibitory	L-DOPA
Serotonin	5-HT ₁ , 5-HT ₂	Excitatory/Inhibitory	5-HTP, SSRIs
Norepinephrine	α, β	Excitatory	Amphetamines, Cocaine
Acetylcholine	Nicotinic, Muscarinic	Excitatory	None
Histamine	H ₁ , H ₂	Excitatory	None
Oxytocin	OxTR	Inhibitory	None
Endocannabinoids	CB ₁ , CB ₂	Inhibitory	None

N Engl J Med 2016 Jan 28;374(4):363-71. doi:10.1056/NEJMr1511480.

19

- Substances take over
- Conditioned behaviors and cues
- Prefrontal cortex loses override power
- Stress/memory can independently activate

Prefrontal cortex (PFC), Anterior cingulate gyrus (ACG), orbitofrontal cortex (OFC), Subgenual cingulate cortex (SCC)

20

Neuroscience cont.

Dopamine Pathways

- Substantia nigra
- VTA
- Mesocortical pathway
- Nigrostriatal pathway
- Tuberoinfundibular pathway
- Mesolimbic pathway

Serotonin Pathways

- Substantia nigra
- Raphe nucleus
- Hypothalamus
- Midbrain
- Pituitary
- Striatal GABA


Functions: Reward (motivation), Pleasure, euphoria, Motor function (fine tuning), Compulsion, Persistence

Effects: Mood, Memory processing, Sleep, Cognition

Brain regions shown: Frontal cortex, Striatum, Hypothalamus, Cerebellum, Dorsal Striatum, Thalamus, Putamen, Nucleus accumbens, Substantia nigra, Ventral tegmental area.

21

Drugs of Abuse and its common effect on VTA and NAC



Nestler EJ: "Is There a Common Molecular Pathway for Addiction?" Nature Neuroscience 8:1445-1449, 2005

22

Understanding Alcohol Use

- Equivalent of the following is considered one drink in the USA
 - 12 oz of regular beer
 - 5 oz of wine
 - 1.5 oz of distilled spirits
- National Institute on Alcohol Abuse and Alcoholism in the elderly recommends the following for healthy people who do NOT take medication
 - One drink a day on average for an elderly man. No more than 2 drinks at any one time.
 - Women should drink even less.
 - People taking medication should further limit use or should not drink at all
 - According to the Dietary Guidelines, adults who do not drink alcohol should not start drinking for any reason.

Source: National Institute of Alcohol Abuse and Alcoholism, www.niaaa.nih.gov and SAHMSA TIP 26
<https://health.gov/our-work/food-sources/2015-2020-dietary-guidelines-guidelines-for-older-adults/>. Accessed 9/30/2020

23

Alcohol Use Continued

- Early onset drinkers:
 - 2/3 of older patients
 - Psychiatric co-occurring are common
 - Severe medical complications secondary to heavy use
- Late onset drinkers:
 - Often triggered by stressful life event
 - More mild cases with fewer medical problems
 - More amenable to treatment

24

Social Factors Contribute to Drinking

- Play an important role in the initiation of AUD(Alcohol Use Disorder)
- Difficult experiences :
 - Loss
 - Physical limitation
 - Isolation
 - Loss of income
 - Loss of occupation

25

Demographics

- Increase the risk of substance use as an older adult:
- Male gender: for EtOH, Cannabis, and tobacco
- Female Gender: For prescription drug use
- Caucasian
- Age (being closer to middle age)
- Less than a college education

26

Physical Factors

- The Following increase the risk of substance use:
- Chronic pain
 - Disabilities and reduce mobility
 - Poor Physical Health
 - Polypharmacy

27

Increased Impact of Alcohol in the Geriatric Population

- Increased Blood Alcohol Concentration because:
 - Decreased lean body mass
 - Decreased total body water
 - Decreased gastric alcohol dehydrogenase
 - Alcohol and drugs more intoxicating in geriatric patients

28

Question

Pt is in ER, BAL is 350mg/d. has no withdrawal symptoms. Has been using Alcohol for 10+ yrs on a daily visit. Has h/o of alcohol withdrawal seizure. Labs are WNL. Poor Social support. He is here is he wants to quit. What will be next step in Management

- A. Give banana bag in ER with Thiamine and Folate
- B. Admit the pt on medical floor with CIWA protocol and Schedule Lorazepam
- C. Admit the pt on observation
- D. Discharge pt
- E. Start pt on Naltrexone

29

Question

Liver biopsy of 65-year-old obese male reveals microvascular fatty globules. He has a 20-year history of heavy alcohol consumption. Lab reveals elevated AST, ALT, low serum albumin, high, globulin and normal bilirubin. What is most likely diagnosis

- A. Alcoholic cirrhosis
- B. Alcoholic liver disease
- C. Wernicke's Korsakoff syndrome
- D. Alcoholic hepatitis
- E. Hepatocellular carcinoma

30

Medical Complications of Alcohol in Geriatric Patients

- Cirrhosis: 60% 1 year death rate > age 60 vs 7% in younger population
- Heart problems (coronary artery disease, and atrial fibrillation)
- Increase in cancers
- Thrombocytopenia
- Neurologic complications (stroke, dementia, Wernicke's encephalopathy)

31

Assessment May Include:

- Skillful Interviewing, willing to ask difficult questions
- Psychiatric evaluation
- Neurological evaluation
- Social Evaluation
- Evaluation of motivation to change
- Functional Evaluation

32

Screening Tools

- Questions about quantity and frequency
 - How many days does the individual drink?
 - Maximum number of drinks on any given occasion
- Instruments:
 - CAGE
 - AUDIT-C
 - MAST-G

33

AUDIT-C

AUDIT-C

Please circle the answer that is correct for you.

1. How often do you have a drink containing alcohol?	2. How many drinks containing alcohol do you have on a typical day when you are drinking?	3. How often do you have six or more drinks on one occasion?	SCORE
Never (0)	1 or 2 (0)	Never (0)	0
Monthly or less (1)	3 or 4 (1)	Less than Monthly (1)	1
Two to four times a month (2)	5 or 6 (2)	Monthly (2)	2
Two to three times per week (3)	7 to 9 (3)	Two to three times per week (3)	3
Four or more times a week (4)	10 or more (4)	Four or more times a week (4)	4
TOTAL SCORE Add the number for each question to get your total score.			0-4

Maximum score is 12. A score of ≥ 4 identifies 86% of men who report drinking above recommended levels or meets criteria for alcohol use disorders. A score of ≥ 2 identifies 84% of women who report hazardous drinking or alcohol use disorders.

34

Protective Factors

- Married
- Supportive, safe living environment
- A provider with knowledge of addiction supervising diverse medications
- Adequate income to meet needs (medical expenses likely to far exceed those of younger adult)
- Annual substance abuse screening including psycho-education. (SAMHSA recommends for 60+)
- Wellness factors including eating, sleeping, exercise, spirituality.
- Linkage to age-specific groups and activities
- Access to transportation

Source: SAMHSA 2015

35

Treatment and Intervention

- Brief Advice
- Brief Interventions
- Facilitates treatment entry and change in behavior
- Referral Management

36

Brief Interventions

- Brief interventions aim to identify a real or potential alcohol problem and motivate an individual to do something about it
- Not designed to treat people with serious dependence

37

Brief Interventions

- Avoid the use of pejorative, labeling words such as "alcoholic" and "abuse"
- The WHO (World Health Organization) has a manual online which outlines brief interventions
- Example Script: "I have looked over the results of the questionnaire you completed a few minutes ago. If you remember, the questions asked about how much alcohol you consume, and whether you have experienced any problems in connection with your drinking. From your answers it appears that you may be at risk of experiencing alcohol-related problems if you continue to drink at your current levels. I would like to take a few minutes to talk with you about it."

Source: WHO http://www.who.int/substance_abuse/activities/brief/

38

Interventions in Geriatric Patients

- Avoid confrontational approaches
- Communicate with empathy in a straightforward, simple manner
- Pay attention to what is important to patients and motivate them (Motivational Interviewing)
- Involve family members or other social support whenever possible

39

Detoxification in Elderly Patients

- **Confusion** (rather than tremor) is an early withdrawal sign
- Duration of withdrawal/hallucinosi s increased
- Rule out Delirium Tremens in confused patients
- Replace electrolytes and nutrients
- Use short acting benzodiazepines (lorazepam, oxazepam)
- Symptomatology monitored with Clinical Institute Withdrawal Assessment for Alcohol (CIWA)
- In the emergency setting, it is imperative to give thiamine, folate and multivitamins early.

Source: LeRoux C, Tang T, Dresler K. Alcohol and Opioid Use Disorder in Older Adults: Neglected and Treatable Illnesses. *Curr Psychiatry Rep*(2016) 18:87

40

A Note About Wernicke's Encephalopathy

- Clinically the classic triad is: ocular findings, cerebellar dysfunction and confusion
- Thiamine needs to be given BEFORE glucose to avoid Wernicke's encephalopathy because glucose depletes thiamine in the body.
- If Wernicke's is suspected, immediate, high dose thiamine is required, some suggest 200mgTID, IV or IM for 3-5 days

Source: Gavin et al. EFNS guidelines for diagnosis, therapy, and prevention of Wernicke encephalopathy. *Eur J Neurol*. 2010;17(12):1408-18.

41

Treatment Strategies

- Age-specific psychosocial approaches are indicated for persons who are not affected with dementia
- Psychotherapy
- Medication management
- Self help groups
- Crisis Management may be needed


42

Pharmacology

- 3 FDA approved medications for AUD:
 - naltrexone
 - acamprosate
 - disulfiram
- Disulfiram is generally not recommended in older adults

43

Naltrexone XR vs Placebo



Garbutt et al, JAMA. 2005;293(13):1617-1625.doi:10.1001/jama.293.13.1617

44

Sedative-Hypnotic Use

- 20% of patients in intermediate care facilities received benzodiazepines (87% as a standing order)
- 41% of psychotropic drug orders in nursing homes are antianxiety agents (mainly benzodiazepines)

Source: Beers M, Avorn J, Soumerai S.B., Everett DE, Shernann DS, Salem S. Psychoactive medication use in intermediate - care facility residents. JAMA 1989; 260: 3016-20.
 Source: Beardsley RS, Larson DB, Burns BJ, Thompson JW, Kamerow DB. Prescribing of psychotropics in elderly nursing home patients. J Am Geriatr Society 1989; 32:7-20.

45

Sedative-Hypnotic Use

- A 2004 study revealed 13% of nursing home residents took benzodiazepines
- Of those residents, 42% did not have an appropriate indication
- This practice was more common in patients that were female, Caucasian, and/or had behavioral disturbance

Source: Stevenson, Decker, Dwyer, Huskamp, Grabowski, Metzger, Mitchell. Antipsychotic and benzodiazepine use among nursing home residents: findings from the 2004 National Nursing Home Survey. American Journal of Geriatric Psychiatry. Dec 16, 2010.

46

Why are sedatives in the elderly concerning?

- Absorption: slower
- Protein binding: elderly patients with low albumin have increased sedation
- Metabolism: slower hepatic metabolism in the elderly; several benzodiazepines have a complicated liver metabolism
- Older patients are at risk of BZD-related harms
 - Fractures
 - Falls
 - Sedation

Take a multifaceted stepwise approach when deprescribing benzodiazepines in older patients. Drugs Ther Perspect 35, 72-76 (2019)

47

Benzodiazepines

- Sedation
- Cerebellar toxicity
- Cognitive impairment
- Psychomotor impairment
- The longer a person is prescribed these medications, the more likely they are to develop misuse
- The longer they are prescribed these medications, the harder it is to taper off

48

How to discontinue Benzodiazepines

- Gradual taper of benzodiazepines is best, especially if medication has been taken chronically, recommendations vary but generally reducing dose by 12-25% every week to month
- Slower clearance of medication attenuates withdrawal symptoms, thus elderly patients *may* report fewer symptoms
- The severity of the distress experienced by a patient during withdrawal is associated with high levels of anxiety, lower educational levels, lower baseline health-related Quality of life, and low levels of social support

49

Guidelines for Prescribing Benzodiazepines

- Prescribe only small dosages
- Prescribe benzodiazepines without active metabolites
- Avoid prescribing benzodiazepines to confused patients or to patients with dementia
- Prescribe benzodiazepines for short periods of time, if at all
- Be aware of potential interaction amongst CNS depressant substances

Saltzman C et al. Clinical Geriatric Psychopharmacology, 3rd Edition, 1998, William & Wilkins 343-355.

50

Prescription drug use

- Elderly patients account for 30% of prescriptions, mainly benzodiazepines and prescription opioids
- One forth of patients in intermediate care facilities are on benzodiazepines
- Rates of benzodiazepine and opioid prescriptions in those 65+ have continued to increase from 2006-2007 to 2014-2015
 - Benzodiazepines from 4.8% to 6.2%
 - Opioids from 5.9% to 10%
 - Benzodiazepines and opioids combined from 1.1% to 2.7%
- The more medications a patient takes, the more likely the medications will be taken improperly

Source: Beers, Avorn et al 1995
Source: Rhee, Coprescribing of Benzodiazepines and Opioids in Older Adults, The Journals of Gerontology December 2019.

51

Question

With statement differentiate the action of cocaine and amphetamines?

- A. Cocaine prevents dopamine reuptake, and amphetamine blocks dopamine and norepinephrine transporters in addition to vesicular monoamine transporters 2 (VMAT-2) inhibition and monoamine oxidase activity inhibition.
- B. Amphetamine prevents dopamine reuptake, but cocaine both slows uptake of dopamine and induces dopamine release.
- C. Both cocaine and amphetamines slow dopamine reuptake and induces dopamine release
- D. Cocaine prevents dopamine reuptake, but amphetamine includes dopamine release
- E. Amphetamine prevents dopamine reuptake, but cocaine includes dopamine release

52

Opioids



53

Opioids

- Opiates: Derived from Opium. Heroin is derived from Opium.
- Opioids: Initially referred to synthetics, now generally refers to synthetic, natural and semi-synthetics

54

Most commonly misused opioid medications

- Oxycodone (OxyContin)
- Oxycodone/acetaminophen (Percocet)
- Hydrocodone (Vicodin)

(Prescription Drugs April 13, 2010)

55

Signs of Opioid Misuse

- Confusion
- Depression
- Delirium
- Insomnia
- Parkinson's-like symptoms
- Weakness or lethargy
- Loss of appetite
- Falls
- Changes in speech; slurring

56

Signs of Opioid Misuse Continued

- Loss of motivation
- Memory loss
- Family or marital discord
- New difficulty with activities of daily living (ADL)
- Drug seeking behavior
- Doctor shopping
- ****Always Check the Physician Monitoring Program (PMP) Before Prescribing! ****

57

Medication Assisted Treatment

- Naltrexone: Oral and Long acting injectable
- Mu-opioid receptor Antagonist
- Paucity of research for IM Naltrexone in older adults

58

Medication Assisted Treatment

- **Buprenorphine:** is a partial mu-opioid receptor agonist with a very high affinity for the mu-opioid receptor
 - Comes in several formulations (SL, Buccal)
 - Must be prescribed by a provider with a DEA X waiver
 - Must be administered when patient is exhibiting symptoms of withdrawal or has already completed withdrawal. In the setting of active intoxication, may precipitate a withdrawal

59

Medication Assisted Treatment

- **Methadone:** A full mu-opioid receptor agonist
 - May only be dispensed from federally regulated opioid treatment programs "methadone clinics"
 - Monitoring required for sedation and respiratory depression
 - May cause CNS depression if used with alcohol, sedatives, hypnotics, or opioids.

60

Naloxone

- Naloxone: Opioid Antagonist
- Patients receiving MAT should have an emergency naloxone kit prescribed
- Consider discussing Naloxone in patients on high dose opioids
- Helpful to engage the patient's family on overdose risk and availability of Naloxone

61

Marijuana

- The medical use of marijuana is legalized in 33 states and DC
- Recreational use is legalized in 11 states
- Use remains federally illegal
- Paucity of research in use in older adults

Mahvan TD, Hilaris ML, Mann A, Brown A, Linn B, Gardner T, Lai B. Marijuana Use in the Elderly: Implications and Considerations. Consult Pharm. 2017 Jun 1;32(6):341-351. doi: 10.4140/TCPh.2017.341. PMID: 28995884.

62

Marijuana

- Acute adverse impact of marijuana use:
 - Anxiety
 - dry mouth
 - tachycardia
 - high blood pressure
 - palpitations
 - wheezing
 - confusion
 - Dizziness

Volkow ND, Baler RD, Compton WM, Weiss SRB. 2014. Adverse health effects of marijuana use. N. Engl. J. Med 370, 2219-2227

63

Marijuana

- Between 2015-2016, prevalence of past year use was 9% in adults ages 50-64 and 2.9% in adults 65 and older
- Based on the National Survey on Drug Use and Health
- Other substance use disorders and misuse of prescription medications were higher in people who used marijuana compared to non-users
- Some concern that older adults who use MJ medically have a higher rate of recreational use too

Han BH, Palamar JJ. Marijuana use by middle-aged and older adults in the United States, 2015-2016. *Drug Alcohol Depend.* 2018 Oct 1;191:374-381

Choi NG, DiNitto DM, Marti N. 2017a. Nonmedical versus medical marijuana use among three age groups of adults: associations with mental and physical health status. *Am. J. Addict.* 26, 697-706.

64

Question

What drugs are FDA approved to treat Tobacco use disorder?

- A. Nortriptyline
- B. Clonidine
- C. Sertraline
- D. Bupropion
- E. Varenicline
- F. All –ab&c
- G. Both d&e

65

Question

What is current prevalence of smoking in United States in Older Adults?

- A. 8%
- B. 16.7%
- C. 17%
- D. 8.2 %

Ref: Cornelius ME, Wang TW, Jamal A, Loretan C, Neff L. Tobacco Product Use Among Adults – United States, 2019.

66

Tobacco Use

- Tobacco use is the leading cause of cancer and death from cancer.
- There is no safe level of tobacco use.
- People who quit smoking, regardless of their age, have gains in life expectancy compared with those who continue to smoke.
- The NCI quitline, 1-877-44U-QUIT (1-877-448-7848), is available Monday through Friday, 9:00 a.m. to 9:00 p.m. ET.

Source: NIH National Cancer Institute. <https://www.cancer.gov/about-cancer/causes-prevention/risk/tobacco>. Accessed on 9/30/2020

67

Thank you

- Questions?

68

References

- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Arlington, VA: American Psychiatric Publishing.
- [Sperio, A., Borge, P., Blazer, D.D., Moore, A.A. Substance abuse among older adults. Clin Geriatr Med. 2014 Aug;30\(5\):829-54. doi: 10.1016/j.cger.2014.04.008. Epub 2014 Jun 12. PMID: 25037298; NCI Thesaurus C127148](#)
- [Dyckhoff, D., Hermal, N., Holbrook, S. Substance use disorders in elderly admissions to an academic psychiatric inpatient service over a 10-year period. Journal of Addiction, Volume 2016, Article ID 4972016](#)
- [Kerns, G.J., Eizenman, J., Frazier, A., et al. The emerging problems of alcohol and substance abuse in late life. J Soc Distress Homeol. 1999;8\(4\):227-230](#)
- National Institute of Alcohol Abuse and Alcoholism. <https://niaaa.nih.gov/alcohol-health/special-populations-co-occurring-disorders/older-adults> accessed on 9/25/17
- [SAMHSA Tip 24: <https://www.samhsa.gov/2k16/2016-09-24-Substance-Abuse-Among-Elderly-Adults-47.pdf>](#)
- <https://health.gov/our-work/food-nutrition/2015-2020-dietary-guidelines/appendix-b/>. Accessed 9/30/2020
- [Bauer, M., Arora, J., Bionardi, S.B., Evans, D.C., Shuman, D.B., Soren, B. Psychotropic medication use in in-home care – care facility residents. JAMA. 1988; 260: 3016-20.](#)
- [Bewick, R.S., Larson, D.B., Burns, B.J., Thompson, J.H., Kasperian, D.B. Prescribing of psychotropics in elderly nursing home patients. J Am Geriatr Society. 1989; 37:37-39.](#)
- [Take a multifaceted stepwise approach when deprescribing benzodiazepines in older patients. Drugs Ther Perspect 35, 72-78 \(2019\)](#)
- [Salmian, C. et al. Clinical Geriatric Psychopharmacology, 3rd Edition, 1999, Wilkins & Wilkins 343-355.](#)
- [Lofthouse, C., Tang, T., Dwyer, K. Alcohol and Opioid Use Disorder in Older Adults: Neglected and Treatable Illnesses. Curr Psychiatry Rep\(2016\) 18:87](#)
- [Gavin, et al. EFNIS guidelines for diagnosis, therapy, and prevention of Wernicke encephalopathy. Eur J Neurol. 2010;17\(12\):1408-15.](#)
- NIH National Cancer Institute. <https://www.cancer.gov/about-cancer/causes-prevention/risk/tobacco>. Accessed on 09/30/2020
- [Steinman, Decker, Dwyer, Hittinger, O'Connell, Mizer, Mitchell. Antipsychotic and benzodiazepine use among nursing home residents: findings from the 2004 National Nursing Home Survey. American Journal of Geriatric Psychiatry. Dec. 18, 2015.](#)
- [Rhee. Coprescribing of Benzodiazepines and Opioids in Older Adults. The Journals of Gerontology December 2019.](#)

69

References

- Han BH, Palamar JJ. Marijuana use by middle-aged and older adults in the United States, 2015-2016. *Drug Alcohol Depend.* 2018 Oct 1;191:374-381. doi: 10.1016/j.drugalcdep.2018.07.026.
- Choi NG, D'Nitto DM, Mars N. 2017a. Nonmedical versus medical marijuana use among three age groups of adults: associations with mental and physical health status. *Am. J. Addict.* 26, 697-706.
- Volkow ND, Baler RD, Compton WM, Weiss SRB. 2014. Adverse health effects of marijuana use. *N. Engl. J. Med.* 370, 2219-2227.
- Kuehse A. Substance Use among Older Adults: An Update on Prevalence, Etiology, Assessment, and Intervention. *Gerontology* 2020;66:249-256. doi: 10.1159/000504363

70

Geriatric Behavioral Health Conference

71

Geriatric Behavioral Health Conference



72
